

isingLenzMC v0.2: Glauber and Metropolis single spin flip dynamics for the Ising Model *

Mehmet Süzen, Ph.D. †

February 28, 2015

The Ising-Lenz Model ¹ appear as one of the land mark systems in statistical physics, as well as in many other fields including computational neuroscience. Because of its simplicity and success in explaining critical phenomenon, Ising Model is routinely used in research and as a teaching concept in statistical mechanics and in standard Monte Carlo methods. There exist analytical solutions for 1D and 2D Ising Model. Still, numerical solutions with MC could provide insights.

isingLenzMC package provides utilities to simulate one dimensional Ising Model with Metropolis and Glauber Monte Carlo with single flip dynamics in periodic boundary conditions ². Computationally intensive parts are written in *low-level language (C)* for efficiency reasons.

1 Ising Model

Consider one dimensional lattice that contains N sites. Each site values can be labelled as $\{s_i\}_{i=1}^N$. In two state version of a lattice, which is an Ising Model, sites can take two values, such as $\{1, -1\}$, corresponding to spin up and spin down states, for example as a model of magnetic material or the state of a neuron.

The total energy, so called Hamiltonian, of the system can be expressed as follows, for short-range parts with interaction strength J :

$$\mathcal{H}(\{s_i\}_{i=1}^N, J, H) = J\left(\sum_{i=1}^{N-1} s_i s_{i+1}\right) + (s_1 s_N) + H \sum_1^N s_i$$

This expression contains two interactions, due to nearest-neighbors (NN) and due to an external field. Coefficients J and H corresponds to these interactions respectively. Note that, additional term in NN interactions $s_1 s_N$ appears due to periodic (cyclic) boundary conditions.

*This package supports following work, *Effective ergodicity in single-spin-flip dynamics*, Phys. Rev. E 90, 032141 (2014) DOI: 10.1103/PhysRevE.90.03214, Mehmet Suezzen

†mehmet.suzen@physics.org

¹E. Ising 1924

²Higher dimensional models may be introduced in future releases.

2 Simulation of the Ising Model

2.1 Single-flip dynamics

One of the common ways to generate dynamics for a lattice system explained in the previous section is changing the value of a randomly chosen site to its opposite value as a dynamical step. This procedure is called single-flip dynamics. An example dynamics for 5 site lattice can be as follows:

$$\begin{aligned} &\{1, -1, 1, 1, -1\} \\ &\{1, -1, 1, -1, -1\} \\ &\{1, 1, 1, -1, -1\} \\ &\{-1, 1, 1, -1, -1\} \end{aligned}$$

Note that, the quality of this kind of dynamics depends on the quality of the random number generator (RNG) we use in selecting flipped site. However, we assume that RNG in use is sufficiently good for this purpose. This matter is beyond the scope of this document, however default RNG in R, Marsenne-Twister is an appropriate choice.

2.2 Transition Probability

The transition probability associated to single spin flip, for Glauber and Metropolis dynamics, can be computed as follows

$$\begin{aligned} p_{Glauber}(\{s_i\}_{i=1}^N) &= \exp(-k\Delta\mathcal{H}) / (1 + \exp(k - \Delta\mathcal{H})) \\ &= 1 / (1 + \exp(k\Delta\mathcal{H})) \\ p_{Metropolis}(\{s_i\}_{i=1}^N) &= \min(1, \exp(-k\Delta\mathcal{H})) \end{aligned}$$

where $k = \frac{1}{k_B T}$ is the Boltzmann factor and $\Delta\mathcal{H}$ is the total energy difference.

2.3 Metropolis Monte Carlo

In metropolis Monte Carlo, the above transition probability is compared with a uniform number between $[0, 1]$ to check the new lattice configuration induced by the spin flip is an acceptable move. Note that the above formulation of the transition probability numerically ensures that transition probability lies in between $[0, 1]$. This procedure mimics importance sampling. This is a special case of Metropolis-Hastings Markov Chain Monte Carlo (MCMC).

3 Utilities

Package provides utilities to perform Metropolis MC for two state 1D N -site lattice, i.e., Ising Model. Functions with suffices with $_R$ are pure R implementations. In this section we document only C based implementations, while functionality is the same.

One can generate a random configuration, perform single flip on the given configuration, compute nearest-neighbour energy and total energy. In the following example we generate 7 sites randomly and perform a single spin flip. We compute energies

```

> require(isingLenzMC)
> set.seed(123456)
> N <- 7
> myInitialConfig <- genConfig1D(N)
> myInitialConfig

[1] 1 1 -1 -1 -1 -1 1

> myNextConfig <- flipConfig1D(myInitialConfig)
> myNextConfig

[1] -1 1 -1 -1 -1 -1 1

> # nearest neighbour energy for initial config
> lattice1DEnergyNN(myInitialConfig)

[1] 3

> # transition probability at J=H=1/kBT=1.0
> transitionProbability1D(1.0, myInitialConfig, myNextConfig, 1.0, 1.0, 1) # Metropolis

[1] 0.002478752

```

It is possible to do the above steps in one go by applying MC move

```

> require(isingLenzMC)
> set.seed(123456)
> N <- 7
> myInitialConfig <- genConfig1D(N)
> myInitialConfig

[1] 1 1 -1 -1 -1 -1 1

> # 1 step Monte Carlo move
> isStep1D(1.0, myInitialConfig, 1.0, 1.0, 1) # Metropolis

$vec
[1] 1 1 -1 -1 -1 -1 1

$accept
[1] 0

```

4 Simulations

4.1 Free Energy

The partition function Z_N can be computed by using the eigenvalues λ_1 and λ_2 of the transfer matrix.

$$Z_N(J, H) = \lambda_1^N + \lambda_2^N$$

For example for 7 sites

```

> Tm <- transferMatrix(1.0, 1.0, 1.0)
> # Free Energy
> log(Tm$values[1]^7 + Tm$values[2]^7)

[1] 14.01997

```

4.2 Magnetisation: Finite Size Effects

The average magnetisation of the 1D ising model is simply defined as the average of the lattice site values in the finite case. However, in theory magnetisation of 1D bulk system ($N \rightarrow \infty$) is analytically known

$$M_{ensemble}(H, T) = \exp(J/k_B T) \sinh(H/k_B T) \left(\exp(2K) \sinh^2(J/k_B T) + \exp(-2J/k_B T) \right)^{-1/2}$$

This is approximately 0.9934346 for $J = H = 1/k_B T = 1.0$. This value represents the ensemble average magnetisation for the bulk system.

Now if we simulate a long enough lattice, we see that simulated magnetisation, time average value, approaches the ensemble average magnetisation value.

```
> require(isingLenzMC)
> set.seed(123456)
> ensembleM <- 0.9934346
> N <- 200
> x <- genConfig1D(N)
> mcData <- isPerform1D(1.0, x, 1.0, 1.0, 10000, ensembleM, 1) # Metropolis
```

a member of *mcData* named list *omegaM* reports so called a fluctuation metric over accepted steps, and it is defined as follows

$$\Omega_M(k) = \left(\left(\sum_{i=1}^k M_{ave}^i \right) - M_{ensemble} \right)^2$$

where M_{ave}^i is the average magnetisation per site at a given accepted step i . It is left as an exercise to see how this fluctuation metric changes with the system size N . The larger the lattice, longer simulation needed to reach to ensemble average.

5 Acknowledgements

Author wishes to thank Dr. Ole Peters, Dr. Richard M. Neumann and Dr. Cornelius Weber for valuable discussions and correspondence.